



## On Topological Properties of Boron Triangular Sheet $BTS(m, n)$ , Borophene Chain $B_{36}(n)$ and Melem Chain $MC(n)$ Nanostructures

Haidar Ali<sup>1,\*</sup>, Abdul Qudair Baig<sup>2</sup>, Muhammad Kashif Shafiq<sup>1</sup>

<sup>1</sup>Department of Mathematics, Government College University, Faisalabad, Pakistan

<sup>2</sup>Department of Mathematics, COMSATS Institute of Information Technology, Attock Campus, Pakistan

**Academic Editor:** Ali Reza Ashrafi

**Abstract.** Topological indices are numerical parameters of a graph which characterize its topology and are usually graph invariant. In QSAR/QSPR study, physico-chemical properties and topological indices such as Randić, atom-bond connectivity ( $ABC$ ) and geometric-arithmetic ( $GA$ ) index are used to predict the bioactivity of chemical compounds. Graph theory has found a considerable use in this area of research. In this paper, we study and derive analytical closed results of general Randić index  $R_\alpha(G)$  with  $\alpha = 1, \frac{1}{2}, -1, -\frac{1}{2}$ , for boron triangular sheet  $BTS(m, n)$ , borophene chain of  $B_{36}(n)$  and melem chain  $MC(n)$ . We also compute the general first Zagreb,  $ABC$ ,  $GA$ ,  $ABC_4$  and  $GA_5$  indices of sheet and chains for the first time and give closed formulas of these degree based indices.

**Keywords.** general Randić index, atom-bond connectivity ( $ABC$ ) index, geometric-arithmetic ( $GA$ ) index, boron triangular, borophene, melem.

### 1 Introduction and preliminary results

Graph theory has provided chemist with a variety of useful tools, such as topological indices. Molecules and molecular compounds are often modeled by molecular graph. A molecular graph is a representation of the structural formula of a chemical compound in terms

\*Corresponding author (Email address: [haidar3830@gmail.com](mailto:haidar3830@gmail.com)).  
DOI: 10.22061/JMNS.2017.705

of graph theory, whose vertices correspond to the atoms of the compound and edges correspond to chemical bonds. *Cheminformatics* is new subject which is a combination of chemistry, mathematics and information science. It studies Quantitative structure-activity (QSAR) and structure-property (QSPR) relationships that are used to predict the biological activities and properties of chemical compounds. In the QSAR /QSPR study, physico-chemical properties and topological indices such as Wiener index, Szeged index, Randić index, Zagreb indices and ABC index are used to predict bioactivity of the chemical compounds.

A graph can be recognized by a numeric number, a polynomial, a sequence of numbers or a matrix. A topological index is a numeric quantity associated with a graph which characterize the topology of graph and is invariant under graph automorphism. There are some major classes of topological indices such as distance based topological indices, degree based topological indices and counting related polynomials and indices of graphs. Among these classes degree based topological indices are of great importance and play a vital role in chemical graph theory and particularly in chemistry. In more precise way, a topological index  $Top(G)$  of a graph, is a number with the property that for every graph  $H$  isomorphic to  $G$ ,  $Top(H) = Top(G)$ . The concept of topological indices came from Wiener [24] while he was working on boiling point of paraffin, named this index as *path number*. Later on, the path number was renamed as *Wiener index* [5].

In this article,  $G$  is considered to be network with vertex set  $V(G)$  and edge set  $E(G)$ ,  $deg(u)$  is the degree of vertex  $u \in V(G)$  and  $S_u = \sum_{v \in N_G(u)} deg(v)$  where  $N_G(u) = \{v \in V(G) \mid uv \in E(G)\}$ . The notations used in this article are mainly taken from books [6, 10].

Let  $G$  be a graph. Then the Wiener index of  $G$  is defined as

$$W(G) = \frac{1}{2} \sum_{(u,v)} d(u,v), \quad (1)$$

where  $(u,v)$  is any ordered pair of vertices in  $G$  and  $d(u,v)$  is  $u-v$  geodesic.

The very first and oldest degree based topological index is *Randić index* [20] denoted by  $R_{-\frac{1}{2}}(G)$  and introduced by Milan Randić and defined as

$$R_{-\frac{1}{2}}(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{deg(u)deg(v)}}. \quad (2)$$

The general Randić index  $R_\alpha(G)$  is the sum of  $(deg(u)deg(v))^\alpha$  over all edges  $e = uv \in E(G)$  defined as

$$R_\alpha(G) = \sum_{uv \in E(G)} (deg(u)deg(v))^\alpha \text{ for } \alpha = 1, \frac{1}{2}, -1, -\frac{1}{2}. \quad (3)$$

An important topological index introduced by Ivan Gutman and Trinajstić is the *Zagreb index* denoted by  $M_1(G)$  and defined as

$$M_1(G) = \sum_{uv \in E(G)} (deg(u) + deg(v)). \quad (4)$$

One of the well-known degree based topological index is *atom-bond connectivity* ( $ABC$ ) index introduced by Estrada *et al.* in [7] and defined as

$$ABC(G) = \sum_{uv \in E(G)} \sqrt{\frac{\deg(u) + \deg(v) - 2}{\deg(u)\deg(v)}}. \quad (5)$$

Another well-known connectivity topological descriptor is *geometric-arithmetic* ( $GA$ ) index which was introduced by Vukičević *et al.* in [23] and defined as

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\deg(u)\deg(v)}}{(\deg(u) + \deg(v))}. \quad (6)$$

Only  $ABC_4$  and  $GA_5$  indices can be computed if we are able to find the edge partition of these interconnection networks based on sum of the degrees of end vertices of each edge in these graphs. The fourth version of  $ABC$  index is introduced by Ghorbani *et al.* [8] and defined as

$$ABC_4(G) = \sum_{uv \in E(G)} \sqrt{\frac{S_u + S_v - 2}{S_u S_v}}. \quad (7)$$

Recently fifth version of  $GA$  index is proposed by Graovac *et al.* [9] and defined as

$$GA_5(G) = \sum_{uv \in E(G)} \frac{2\sqrt{S_u S_v}}{(S_u + S_v)}. \quad (8)$$

The general Randić index for  $\alpha = 1$  is the second Zagreb index for any graph  $G$ .

## 2 Main Results

We study the general Randić, first Zagreb,  $ABC$ ,  $GA$ ,  $ABC_4$  and  $GA_5$  indices and give closed formulae of these indices for boron triangular sheet  $BTS(m, n)$ , borophene chain of  $B_{36}(n)$  and melem chain  $MC(n)$ . Imran *et al.* studied various degree based topological indices for various networks like silicates, hexagonal, honeycomb and oxide in [12]. Nowadays there is an extensive research activity on  $ABC$  and  $GA$  indices and their variants, for further study of topological indices of various graph families see, [1–4, 13–19, 21, 22].

### 2.1 Results for $BTS(m, n)$ , $B_{36}(n)$ and $MC(n)$ Nanostructures

In this paper, we calculate certain degree based topological indices of boron triangular sheet  $BTS(m, n)$ , borophene chain of  $B_{36}(N)$  and melem chain  $MC(n)$  nanostructures. We compute general Randić  $R_\alpha(G)$  with  $\alpha = \{1, -1, \frac{1}{2}, -\frac{1}{2}\}$ ,  $ABC$ ,  $GA$ ,  $ABC_4$  and  $GA_5$  indices for  $BTS(m, n)$ ,  $B_{36}(n)$  and  $MC(n)$  nanostructures.




Figure 1. Boron triangular sheet (BTS(4, 4))




Figure 2. Borophene chain ( $B_{36}(n)(3)$ )

**Theorem 2.1.** Consider the boron triangular sheet  $BTS(m, n)$  for  $m = n \geq 3$ . Then

$$R_{\alpha}(BTS(m, n)) = \begin{cases} -2(7m - 108mn + 7(2 + n)), & \alpha = 1; \\ 12 + 8\sqrt{3} + 4(-4 + m + n) + \\ (4\sqrt{6} + 2\sqrt{15} + 3\sqrt{30})(-2 + m + n) + \\ 3\sqrt{2}(4 + m + n) - 36(-1 + m - mn + n), & \alpha = \frac{1}{2}; \\ \frac{1}{720}(204 + 193m + 120mn + 193n), & \alpha = -1; \\ \frac{1}{60}(80 + 40\sqrt{3} + 15(-4 + m + n) + \\ (10\sqrt{6} + 8\sqrt{5} + 6\sqrt{30})(-2 + m + n) + \\ 10\sqrt{2}(4 + m + n) - 60(-1 + m - mn + n)), & \alpha = -\frac{1}{2}. \end{cases}$$

*Proof.* Let  $G \cong BTS(m, n)$  be the boron triangular sheet. The boron triangular sheet  $BTS(m, n)$  has  $m + n + 4$  vertices of degree 3,  $m + n - 2$  vertices of degree 4,  $m + n - 2$  vertices of degree 5 and  $2mn - m - n + 1$  vertices of degree 6. The edge set of  $BTS(m, n)$  is divided into eight partitions based on the degree of end vertices. The first edge partition  $E_1(BTS(m, n))$  contains 4 edges  $uv$ , where  $deg(u) = deg(v) = 3$ . The second edge partition  $E_2(BTS(m, n))$




Figure 3. Melem chain (MC(4))

Table 1. Edge partition of boron triangular sheet  $BTS(m, n)$  based on degrees of end vertices of each edge.

$(d_u, d_v), (uv \in E(G))$	Number of edges
(3,3)	4
(3,4)	4
(3,5)	$2(m + n - 2)$
(3,6)	$m + n + 4$
(4,4)	$m + n - 4$
(4,6)	$2(m + n - 2)$
(5,6)	$3(m + n - 2)$
(6,6)	$6(mn - (m + n) + 1)$

contains 4 edges  $uv$ , where  $deg(u) = 3$  and  $deg(v) = 4$ . The third edge partition  $E_3(BTS(m, n))$  contains  $2m + 2n - 4$  edges  $uv$ , where  $deg(u) = 3$  and  $deg(v) = 5$ . The fourth edge partition  $E_4(BTS(m, n))$  contains  $m + n + 4$  edges  $uv$ , where  $deg(u) = 3$  and  $deg(v) = 6$ . The fifth edge partition  $E_5(BTS(m, n))$  contains  $m + n - 4$  edges  $uv$ , where  $deg(u) = deg(v) = 4$ . The sixth edge partition  $E_6(BTS(m, n))$  contains  $2m + 2n - 4$  edges  $uv$ , where  $deg(u) = 4$  and  $deg(v) = 6$ . The seventh edge partition  $E_7(BTS(m, n))$  contains  $3m + 3n - 6$  edges  $uv$ , where  $deg(u) = 5$  and  $deg(v) = 6$  and the eighth edge partition  $E_8(BTS(m, n))$  contains  $6mn - 6m - 6n + 6$  edges  $uv$ , where  $deg(u) = deg(v) = 6$ . Table 1 shows such an edge partition of  $BTS(m, n)$ . Thus from (3) it follows that

$$R_\alpha(G) = \sum_{uv \in E(G)} (deg(u) \cdot deg(v))^\alpha.$$

Now, we apply the formula of  $R_\alpha(G)$  for  $\alpha = 1$

$$R_1(G) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} deg(u) \cdot deg(v).$$

By using edge partition given in Table 1, we get

$$\begin{aligned} R_1(G) &= 9|E_1(BTS(m, n))| + 12|E_2(BTS(m, n))| + 15|E_3(BTS(m, n))| + 18|E_4(BTS(m, n))| \\ &\quad + 16|E_5(BTS(m, n))| + 24|E_6(BTS(m, n))| + 30|E_7(BTS(m, n))| + 36|E_8(BTS(m, n))| \\ &= -2(7m - 108mn + 7(2 + n)). \end{aligned}$$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = \frac{1}{2}$

$$R_{\frac{1}{2}}(G) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} \sqrt{deg(u) \cdot deg(v)}.$$

By using edge partition given in Table 1, we get

$$\begin{aligned}
 R_{\frac{1}{2}}(G) &= 3|E_1(BTS(m,n))| + 2\sqrt{3}|E_2(BTS(m,n))| + \sqrt{15}|E_3(BTS(m,n))| \\
 &\quad + 3\sqrt{2}|E_4(BTS(m,n))| + 4|E_5(BTS(m,n))| + 2\sqrt{6}|E_6(BTS(m,n))| \\
 &\quad + \sqrt{30}|E_7(BTS(m,n))| + 6|E_8(BTS(m,n))| \\
 &= 12 + 8\sqrt{3} + 4(-4 + m + n) + (4\sqrt{6} + 2\sqrt{15} + 3\sqrt{30})(-2 + m + n) \\
 &\quad + 3\sqrt{2}(4 + m + n) - 36(-1 + m - mn + n).
 \end{aligned}$$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = -1$ . Then we have

$$\begin{aligned}
 R_{-1}(G) &= \sum_{j=1}^8 \sum_{uv \in E_j(G)} \frac{1}{deg(u) \cdot deg(v)} \\
 &= \frac{1}{9}|E_1(BTS(m,n))| + \frac{1}{12}|E_2(BTS(m,n))| + \frac{1}{15}|E_3(BTS(m,n))| \\
 &\quad + \frac{1}{18}|E_4(BTS(m,n))| + \frac{1}{16}|E_5(BTS(m,n))| + \frac{1}{24}|E_6(BTS(m,n))| \\
 &\quad + \frac{1}{30}|E_7(BTS(m,n))| + \frac{1}{36}|E_8(BTS(m,n))| \\
 &= \frac{1}{720}(204 + 193m + 120mn + 193n).
 \end{aligned}$$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = -\frac{1}{2}$ . Then we have

$$\begin{aligned}
 R_{-\frac{1}{2}}(G) &= \sum_{j=1}^8 \sum_{uv \in E_j(G)} \frac{1}{\sqrt{deg(u) \cdot deg(v)}} \\
 &= \frac{1}{3}|E_1(BTS(m,n))| + \frac{\sqrt{3}}{6}|E_2(BTS(m,n))| + \frac{1}{\sqrt{15}}|E_3(BTS(m,n))| \\
 &\quad + \frac{\sqrt{2}}{6}|E_4(BTS(m,n))| + \frac{1}{4}|E_5(BTS(m,n))| + \frac{\sqrt{6}}{12}|E_6(BTS(m,n))| \\
 &\quad + \frac{1}{\sqrt{30}}|E_7(BTS(m,n))| + \frac{1}{6}|E_8(BTS(m,n))| \\
 &= \frac{1}{60}(80 + 40\sqrt{3} + 15(-4 + m + n) + (10\sqrt{6} + 8\sqrt{5} + 6\sqrt{30})(-2 + m + n) \\
 &\quad + 10\sqrt{2}(4 + m + n) - 60(-1 + m - mn + n)).
 \end{aligned}$$

□

In the following, we compute first Zagreb index of boron triangular sheet  $BTS(m,n)$ .

**Theorem 2.2.** For boron triangular sheet  $G \cong BTS(m,n)$  for  $m = n \geq 3$ , We have

$$M_1(BTS(m,n)) = 2(-5 + 7m + 36mn + 7n).$$

*Proof.* Let  $G$  be the boron triangular sheet  $BTS(m, n)$ . By using edge partition from Table 1, the result follows. From (4) we have

$$\begin{aligned} M_1(BTS(m, n)) &= \sum_{uv \in E(G)} (deg(u) + deg(v)) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} (deg(u) + deg(v)) \\ &= 6|E_1(BTS(m, n))| + 7|E_2(BTS(m, n))| + 8|E_3(BTS(m, n))| \\ &\quad + 9|E_4(BTS(m, n))| + 8|E_5(BTS(m, n))| + 10|E_6(BTS(m, n))| \\ &\quad + 11|E_7(BTS(m, n))| + 12|E_8(BTS(m, n))|. \end{aligned}$$

By doing some calculation, we get  $M_1(BTS(m, n)) = 2(-5 + 7m + 36mn + 7n)$ . □

Now, we compute  $ABC$  and  $GA$  indices of boron triangular sheet  $BTS(m, n)$ .

**Theorem 2.3.** Let  $G \cong BTS(m, n)$  be the boron triangular sheet, for  $m = n \geq 3$ , then

$$\begin{aligned} ABC(G) &= \frac{1}{60}(160 + 40\sqrt{15} + 15\sqrt{6}(-4 + m + n) \\ &\quad + (40\sqrt{3} + 24\sqrt{10} + 18\sqrt{30})(-2 + m + n) \\ &\quad + 10\sqrt{14}(4 + m + n) - 60\sqrt{10}(-1 + m - mn + n)), \\ GA(G) &= 6 + \frac{16}{7}\sqrt{3} - 5m + 6mn - 5n \\ &\quad + \left(\frac{4}{5}\sqrt{6} + \frac{\sqrt{15}}{2} + \frac{6}{11}\sqrt{30}(-2 + m + n) + \frac{2}{3}\sqrt{2}(4 + m + n)\right). \end{aligned}$$

*Proof.* By using edge partition given in Table 1, we get the result. From (5) it follows that

$$\begin{aligned} ABC(G) &= \sum_{uv \in E(G)} \sqrt{\frac{deg(u) + deg(v) - 2}{deg(u) \cdot deg(v)}} = \sum_{j=1}^8 \sum_{uv \in E_j(G)} \sqrt{\frac{deg(u) + deg(v) - 2}{deg(u) \cdot deg(v)}} \\ &= \frac{2}{3}|E_1(BTS(m, n))| + \frac{\sqrt{15}}{6}|E_2(BTS(m, n))| + \frac{\sqrt{10}}{5}|E_3(BTS(m, n))| \\ &\quad + \frac{\sqrt{14}}{6}|E_4(BTS(m, n))| + \frac{\sqrt{6}}{4}|E_5(BTS(m, n))| + \frac{\sqrt{3}}{3}|E_6(BTS(m, n))| \\ &\quad + \frac{\sqrt{30}}{10}|E_7(BTS(m, n))| + \frac{\sqrt{10}}{6}|E_8(BTS(m, n))|. \end{aligned}$$

By doing some calculation, we get

$$\begin{aligned} ABC(G) &= \frac{1}{60}(160 + 40\sqrt{15} + 15\sqrt{6}(-4 + m + n) \\ &\quad + (40\sqrt{3} + 24\sqrt{10} + 18\sqrt{30})(-2 + m + n) + 10\sqrt{14}(4 + m + n) \\ &\quad - 60\sqrt{10}(-1 + m - mn + n)), \end{aligned}$$

and from (6) we get

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\deg(u)\deg(v)}}{(\deg(u) + \deg(v))} \sum_{j=1}^8 \sum_{uv \in E_j(G)} \frac{2\sqrt{\deg(u)\deg(v)}}{(\deg(u) + \deg(v))}.$$

Then we have

$$\begin{aligned} GA(G) &= |E_1(BTS(m,n))| + \frac{4}{7}\sqrt{3}|E_2(BTS(m,n))| + \frac{\sqrt{15}}{4}|E_3(BTS(m,n))| \\ &+ \frac{2}{3}\sqrt{2}|E_4(BTS(m,n))| + |E_5(BTS(m,n))| + \frac{2}{5}\sqrt{6}|E_6(BTS(m,n))| \\ &+ \frac{2}{11}\sqrt{30}|E_7(BTS(m,n))| + |E_8(BTS(m,n))|. \end{aligned}$$

By doing some calculation, we get

$$\begin{aligned} GA(G) &= 6 + \frac{16}{7}\sqrt{3} - 5m + 6mn - 5n \\ &+ \left(\frac{4}{5}\sqrt{6} + \frac{\sqrt{15}}{2} + \frac{6}{11}\sqrt{30}(-2 + m + n) + \frac{2}{3}\sqrt{2}(4 + m + n)\right). \end{aligned}$$

□

Now, we compute  $ABC_4$  and  $GA_5$  indices of boron triangular sheet  $BTS(m,n)$ . Let us consider an edge partition based on degree sum of neighbors of end vertices. Then the edge set  $E(BTS(m,n))$  can be divided into twenty four edge partitions  $E_j(BTS(m,n)), 9 \leq j \leq 32$ , where the edge partition  $E_9(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 13$  and  $S_v = 14$ , the edge partition  $E_{10}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 13$  and  $S_v = 19$ , the edge partition  $E_{11}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 13$  and  $S_v = 27$ , the edge partition  $E_{12}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 14$  and  $S_v = 24$ , the edge partition  $E_{13}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 14$  and  $S_v = 27$ , the edge partition  $E_{14}(BTS(m,n))$  contains  $2m + 2n - 8$  edges  $uv$  with  $S_u = 16$  and  $S_v = 24$ , the edge partition  $E_{15}(BTS(m,n))$  contains  $m + n - 4$  edges  $uv$  with  $S_u = 16$  and  $S_v = 31$ ,  $E_{16}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 19$  and  $S_v = 20$ ,  $E_{17}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 19$  and  $S_v = 27$ ,  $E_{18}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 19$  and  $S_v = 32$ ,  $E_{19}(BTS(m,n))$  contains  $m + n - 8$  edges  $uv$  with  $S_u = S_v = 20$ ,  $E_{20}(BTS(m,n))$  contains  $2m + 2n - 12$  edges  $uv$  with  $S_u = 20$  and  $S_v = 32$ ,  $E_{21}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 24$  and  $S_v = 27$ ,  $E_{22}(BTS(m,n))$  contains  $2m + 2n - 8$  edges  $uv$  with  $S_u = 24$  and  $S_v = 31$ ,  $E_{23}(BTS(m,n))$  contains  $m + n - 2$  edges  $uv$  with  $S_u = 24$  and  $S_v = 35$ ,  $E_{24}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 27$  and  $S_v = 32$ ,  $E_{25}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 27$  and  $S_v = 35$ ,  $E_{26}(BTS(m,n))$  contains  $2m + 2n - 8$  edges  $uv$  with  $S_u = 31$  and  $S_v = 35$ ,  $E_{27}(BTS(m,n))$  contains  $m + n - 4$  edges  $uv$  with  $S_u = 31$  and  $S_v = 36$ ,  $E_{28}(BTS(m,n))$  contains  $m + n - 6$  edges  $uv$  with  $S_u = S_v = 32$ ,  $E_{29}(BTS(m,n))$  contains 4 edges  $uv$  with  $S_u = 32$  and  $S_v = 35$ ,  $E_{30}(BTS(m,n))$  contains  $2m + 2n - 12$  edges  $uv$  with  $S_u = 32$  and  $S_v = 36$ ,  $E_{31}(BTS(m,n))$  contains  $3m + 3n - 10$  edges  $uv$  with  $S_u = 35$  and  $S_v = 36$  and  $E_{32}(BTS(m,n))$  contains  $6mn - 15m - 15n + 34$  edges  $uv$  with  $S_u = S_v = 36$ .



**Theorem 2.4.** Let  $G \cong BTS(m, n)$  be the boron triangular sheet, for  $m = n \geq 5$ , then

$$\begin{aligned}
 ABC_4(G) &= 10\sqrt{\frac{2}{91}} + 4\sqrt{\frac{30}{247}} + \frac{8}{3}\sqrt{\frac{11}{57}} + 2\sqrt{\frac{37}{95}} + 2\sqrt{\frac{3}{7}} + \sqrt{\frac{13}{14}} + \frac{7}{9}\sqrt{2} + \frac{2}{3}\sqrt{\frac{26}{7}} \\
 &+ \frac{8}{3\sqrt{7}} + \frac{1}{3}\sqrt{\frac{19}{2}} + \frac{7}{\sqrt{38}} + \frac{152}{3\sqrt{39}} + \frac{1}{18}\sqrt{\frac{35}{2}}(34 - 15m + 6mn - 15n) \\
 &+ \frac{1}{10}\sqrt{\frac{19}{2}}(-8 + m + n) + \left(\frac{1}{4}\sqrt{\frac{11}{3}} + \frac{1}{4}\sqrt{5} + \frac{1}{16}\sqrt{\frac{31}{2}}\right)(-6 + m + n) \\
 &+ \left(\frac{3}{4}\sqrt{\frac{5}{31}} + \sqrt{\frac{53}{186}} + \frac{1}{6}\sqrt{\frac{65}{31}} + \frac{1}{4}\sqrt{\frac{19}{3}} + \frac{16}{\sqrt{1085}}\right)(-4 + m + n) \\
 &+ \frac{1}{2}\sqrt{\frac{19}{70}}(-2 + m + n) + \frac{1}{2}\sqrt{\frac{23}{105}}(-10 + 3m + 3n), \\
 GA_5(G) &= 20 + \frac{48}{17}\sqrt{2} + \frac{96}{59}\sqrt{6} + \frac{16}{19}\sqrt{21} + \frac{32}{51}\sqrt{38} + \frac{3}{5}\sqrt{39} + \frac{24}{41}\sqrt{42} \\
 &+ \frac{12}{23}\sqrt{57} + \frac{32}{67}\sqrt{70} + \frac{16}{39}\sqrt{95} + \frac{12}{31}\sqrt{105} + \frac{8}{27}\sqrt{182} + \frac{1}{4}\sqrt{247} \\
 &- 13m + 6mn - 13n + \left(\frac{24}{17}\sqrt{2} + \frac{8}{13}\sqrt{10}\right)(-6 + m + n) \\
 &+ \left(\frac{4}{5}\sqrt{6} + \frac{1100}{3149}\sqrt{31} + \frac{8}{55}\sqrt{186} + \frac{2}{33}\sqrt{1085}\right)(-4 + m + n) \\
 &+ \frac{4}{59}\sqrt{210}(-2 + m + n) + \frac{12}{71}\sqrt{35}(-10 + 3m + 3n).
 \end{aligned}$$

*Proof.* By using edge partition given in Table 2, we get the result. From (7) it follows that

$$\begin{aligned}
 ABC_4(G) &= \sum_{uv \in E(G)} \sqrt{\frac{S_u + S_v - 2}{S_u S_v}} = \sum_{j=9}^{32} \sum_{uv \in E_j(G)} \sqrt{\frac{S_u + S_v - 2}{S_u S_v}} \\
 &= \frac{5}{\sqrt{182}}|E_9(BTS(m, n))| + \sqrt{\frac{30}{247}}|E_{10}(BTS(m, n))| + \frac{1}{3}\sqrt{\frac{38}{39}}|E_{11}(BTS(m, n))| \\
 &+ \frac{\sqrt{21}}{14}|E_{12}(BTS(m, n))| + \frac{\sqrt{182}}{42}|E_{13}(BTS(m, n))| + \frac{\sqrt{57}}{24}|E_{14}(BTS(m, n))| \\
 &+ \frac{3}{4}\sqrt{\frac{5}{31}}|E_{15}(BTS(m, n))| + \frac{1}{2}\sqrt{\frac{37}{95}}|E_{16}(BTS(m, n))| + \frac{2}{3}\sqrt{\frac{11}{57}}|E_{17}(BTS(m, n))| \\
 &+ \frac{1}{4}\frac{7}{\sqrt{38}}|E_{18}(BTS(m, n))| + \frac{\sqrt{38}}{20}|E_{19}(BTS(m, n))| + \frac{\sqrt{5}}{8}|E_{20}(BTS(m, n))| \\
 &+ \frac{7}{36}\sqrt{2}|E_{21}(BTS(m, n))| + \frac{1}{2}\sqrt{\frac{53}{186}}|E_{22}(BTS(m, n))| + \frac{3}{2}\sqrt{\frac{6}{210}}|E_{23}(BTS(m, n))| \\
 &+ \frac{\sqrt{38}}{24}|E_{24}(BTS(m, n))| + \frac{2}{21}\sqrt{7}|E_{25}(BTS(m, n))| + \frac{8}{\sqrt{1085}}|E_{26}(BTS(m, n))| \\
 &+ \frac{1}{6}\sqrt{\frac{65}{31}}|E_{27}(BTS(m, n))| + \frac{\sqrt{62}}{32}|E_{28}(BTS(m, n))| + \frac{\sqrt{182}}{56}|E_{29}(BTS(m, n))|
 \end{aligned}$$

$$+ \frac{\sqrt{33}}{24} |E_{30}(BTS(m, n))| + \frac{1}{6} \sqrt{\frac{69}{35}} |E_{31}(BTS(m, n))| + \frac{\sqrt{70}}{36} |E_{32}(BTS(m, n))|.$$

Thus, we have

$$\begin{aligned} ABC_4(G) = & 10\sqrt{\frac{2}{91}} + 4\sqrt{\frac{30}{247}} + \frac{8}{3}\sqrt{\frac{11}{57}} + 2\sqrt{\frac{37}{95}} + 2\sqrt{\frac{3}{7}} + \sqrt{\frac{13}{14}} + \frac{7}{9}\sqrt{2} + \frac{2}{3}\sqrt{\frac{26}{7}} \\ & + \frac{8}{3\sqrt{7}} + \frac{1}{3}\sqrt{\frac{19}{2}} + \frac{7}{\sqrt{38}} + \frac{152}{3\sqrt{39}} + \frac{1}{18}\sqrt{\frac{35}{2}}(34 - 15m + 6mn - 15n) \\ & + \frac{1}{10}\sqrt{\frac{19}{2}}(-8 + m + n) + \left(\frac{1}{4}\sqrt{\frac{11}{3}} + \frac{1}{4}\sqrt{5} + \frac{1}{16}\sqrt{\frac{31}{2}}\right)(-6 + m + n) \\ & + \left(\frac{3}{4}\sqrt{\frac{5}{31}} + \sqrt{\frac{53}{186}} + \frac{1}{6}\sqrt{\frac{65}{31}} + \frac{1}{4}\sqrt{\frac{19}{3}} + \frac{16}{\sqrt{1085}}\right)(-4 + m + n) \\ & + \frac{1}{2}\sqrt{\frac{19}{70}}(-2 + m + n) + \frac{1}{2}\sqrt{\frac{23}{105}}(-10 + 3m + 3n). \end{aligned}$$

From (8) we get

$$GA_5(G) = \sum_{uv \in E(G)} \frac{2\sqrt{S_u S_v}}{(S_u + S_v)} = \sum_{j=9}^{32} \sum_{uv \in E_j(G)} \frac{2\sqrt{S_u S_v}}{(S_u + S_v)}.$$

Then,

$$\begin{aligned} GA_5(G) = & 2\frac{\sqrt{182}}{27} |E_9(BTS(m, n))| + \frac{\sqrt{247}}{16} |E_{10}(BTS(m, n))| + 3\frac{\sqrt{39}}{20} |E_{11}(BTS(m, n))| \\ & + 4\frac{\sqrt{21}}{19} |E_{12}(BTS(m, n))| + 6\frac{\sqrt{42}}{41} |E_{13}(BTS(m, n))| + 2\frac{\sqrt{6}}{5} |E_{14}(BTS(m, n))| \\ & + 8\frac{\sqrt{31}}{47} |E_{15}(BTS(m, n))| + 4\frac{\sqrt{95}}{39} |E_{16}(BTS(m, n))| + 3\frac{\sqrt{57}}{23} |E_{17}(BTS(m, n))| \\ & + 8\frac{\sqrt{38}}{51} |E_{18}(BTS(m, n))| + |E_{19}(BTS(m, n))| + 4\frac{\sqrt{10}}{13} |E_{20}(BTS(m, n))| \\ & + 12\frac{\sqrt{2}}{17} |E_{21}(BTS(m, n))| + 4\frac{\sqrt{186}}{55} |E_{22}(BTS(m, n))| + 4\frac{\sqrt{210}}{59} |E_{23}(BTS(m, n))| \\ & + 24\frac{\sqrt{6}}{59} |E_{24}(BTS(m, n))| + 3\frac{\sqrt{105}}{31} |E_{25}(BTS(m, n))| + \frac{\sqrt{1085}}{33} |E_{26}(BTS(m, n))| \\ & + 12\frac{\sqrt{31}}{67} |E_{27}(BTS(m, n))| + |E_{28}(BTS(m, n))| + 8\frac{\sqrt{70}}{67} |E_{29}(BTS(m, n))| \\ & + 12\frac{\sqrt{2}}{17} |E_{30}(BTS(m, n))| + 12\frac{\sqrt{35}}{71} |E_{31}(BTS(m, n))| + |E_{32}(BTS(m, n))|. \end{aligned}$$

Thus, we have

$$\begin{aligned}
 GA_5(G) &= 20 + \frac{48}{17}\sqrt{2} + \frac{96}{59}\sqrt{6} + \frac{16}{19}\sqrt{21} + \frac{32}{51}\sqrt{38} + \frac{3}{5}\sqrt{39} + \frac{24}{41}\sqrt{42} + \frac{12}{23}\sqrt{57} + \frac{32}{67}\sqrt{70} \\
 &+ \frac{16}{39}\sqrt{95} + \frac{12}{31}\sqrt{105} + \frac{8}{27}\sqrt{182} + \frac{1}{4}\sqrt{247} - 13m + 6mn - 13n \\
 &+ \left(\frac{24}{17}\sqrt{2} + \frac{8}{13}\sqrt{10}\right)(-6 + m + n) \\
 &+ \left(\frac{4}{5}\sqrt{6} + \frac{1100}{3149}\sqrt{31} + \frac{8}{55}\sqrt{186} + \frac{2}{33}\sqrt{1085}\right)(-4 + m + n) \\
 &+ \frac{4}{59}\sqrt{210}(-2 + m + n) + \frac{12}{71}\sqrt{35}(-10 + 3m + 3n).
 \end{aligned}$$

□

Chemical engineers have determined that a unique arrangement of 36 boron-atoms in a flat disc with a hexagonal hole in the middle may be preferred building blocks for borophene. A 36-atom cluster of boron, left, arranged as a flat disc with a hexagonal hole in the middle, fix the theoretical requirements for making a one-atom-thick boron chain, right, a theoretical nanomaterial dubbed borophene. A borophene chain  $B_{36}(n)$  for  $n \geq 2$  has order  $32n + 4$  and size  $81n + 3$ .

Now, we calculate certain degree based topological indices of borophene chain  $B_{36}(n)$  of dimension  $n$ . In the coming theorems we compute general Randić index  $R_\alpha(G)$  with  $\alpha = \{1, -1, \frac{1}{2}, -\frac{1}{2}\}$ ,  $ABC$ ,  $GA$ ,  $ABC_4$  and  $GA_5$  of  $B_{36}(n)$ .

**Theorem 2.5.** Consider the borophene chain  $B_{36}(n)$  for  $n \geq 2$ . Then

$$R_\alpha(B_{36}(n)) = \begin{cases} 6(-32 + 373n), & \alpha = 1; \\ 8 + 8\sqrt{5}(-1 + n) + 46n + (6\sqrt{2} + 8\sqrt{3})(2 + n) + \\ 16\sqrt{6}(1 + 2n) + 6\sqrt{30}(-1 + 4n) + 18(-3 + 7n), & \alpha = \frac{1}{2}; \\ \frac{1}{1800}(1255 + 5732n), & \alpha = -1; \\ \frac{1}{30}(-30 + 20\sqrt{2} + 40\sqrt{3} - 12\sqrt{5} + 20\sqrt{6} - 6\sqrt{30} + \\ (171 + 10\sqrt{2} + 20\sqrt{3} + 12\sqrt{3} + 40\sqrt{6} + 24\sqrt{30})n), & \alpha = -\frac{1}{2}. \end{cases}$$

*Proof.* Let  $G$  be the borophene chain  $B_{36}(n)$ . The borophene chain  $B_{36}(n)$  has  $2n + 4$  vertices of degree 3,  $8n + 4$  vertices of degree 4,  $8n - 2$  vertices of degree 5 and  $14n - 2$  vertices of degree 6. The edge set of  $B_{36}(n)$  is divided into eight partitions based on the degree of end vertices. The first edge partition  $E_1(B_{36}(n))$  contains  $4n + 8$  edges  $uv$ , where  $deg(u) = 3$  and  $deg(v) = 4$ . The second edge partition  $E_2(B_{36}(n))$  contains  $2n + 4$  edges  $uv$ , where  $deg(u) = 3$  and  $deg(v) = 6$ . The third edge partition  $E_3(B_{36}(n))$  contains  $4n + 2$  edges  $uv$ , where  $deg(u) = deg(v) = 4$ . The fourth edge partition  $E_4(B_{36}(n))$  contains  $4n - 4$  edges  $uv$ , where  $deg(u) = 4$  and  $deg(v) = 5$ . The fifth edge partition  $E_5(B_{36}(n))$  contains  $16n + 8$  edges  $uv$ , where  $deg(u) = 4$  and  $deg(v) = 6$ . The sixth edge partition  $E_6(B_{36}(n))$  contains  $6n$  edges  $uv$ , where  $deg(u) = deg(v) = 5$ . The seventh edge partition  $E_7(B_{36}(n))$  contains  $24n - 6$  edges

Table 2. Edge partition of boron triangular sheet  $BTS(m, n)$  based on degrees sum of end vertices of each edge.

$(S_u, S_v), uv \in E(G)$	Number of edges	$(S_u, S_v), uv \in E(G)$	Number of edges
(13,14)	4	(24,27)	4
(13,19)	4	(24,31)	$2m + 2n - 8$
(13,27)	4	(24,35)	$m + n - 2$
(14,24)	4	(27,32)	4
(14,27)	4	(27,35)	4
(16,24)	$2m + 2n - 8$	(31,35)	$2m + 2n - 8$
(16,31)	$m + n - 4$	(31,36)	$m + n - 4$
(19,20)	4	(32,32)	$m + n - 6$
(19,27)	4	(32,35)	4
(19,32)	4	(32,36)	$2m + 2n - 12$
(20,20)	$m + n - 8$	(35,36)	$3m + 3n - 10$
(20,32)	$2m + 2n - 12$	(36,36)	$6mn - 15(m + n) + 34$

Table 3. Edge partition of borophene chain  $B_{36}(n)$  based on degrees of end vertices of each edge.

$(d_u, d_v), uv \in E(G)$	Number of edges
(3,4)	$4n + 8$
(3,6)	$2n + 4$
(4,4)	$4n + 2$
(4,5)	$4n - 4$
(4,6)	$16n + 8$
(5,5)	$6n$
(5,6)	$24n - 6$
(6,6)	$21n - 9$

$uv$ , where  $deg(u) = 5$  and  $deg(v) = 6$ . The eight edge partition  $E_8(B_{36}(n))$  contains  $21n - 9$  edges  $uv$ , where  $deg(u) = deg(v) = 6$ . Table 3 shows such an edge partition of  $B_{36}(n)$ . Thus from (3) it follows that

$$R_\alpha(G) = \sum_{uv \in E(G)} (deg(u)deg(v))^\alpha.$$

Now we apply the formula of  $R_\alpha(G)$  for  $\alpha = 1$

$$R_1(G) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} deg(u) \cdot deg(v).$$

By using edge partition given in Table 3, we get

$$R_1(G) = 12|E_1(B_{36}(n))| + 18|E_2(B_{36}(n))| + 16|E_3(B_{36}(n))| + 20|E_4(B_{36}(n))| + 24|E_5(B_{36}(n))| + 25|E_6(B_{36}(n))| + 30|E_7(B_{36}(n))| + 36|E_8(B_{36}(n))|.$$

Then  $R_1(G) = 6(-32 + 373n)$ . We apply the formula of  $R_\alpha(G)$  for  $\alpha = \frac{1}{2}$

$$R_{\frac{1}{2}}(G) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} \sqrt{\deg(u) \cdot \deg(v)}.$$

By using edge partition given in Table 3, we get

$$R_{\frac{1}{2}}(G) = 2\sqrt{3}|E_1(B_{36}(n))| + 3\sqrt{2}|E_2(B_{36}(n))| + 4|E_3(B_{36}(n))| + 2\sqrt{5}|E_4(B_{36}(n))| \\ + 2\sqrt{6}|E_5(B_{36}(n))| + 5|E_6(B_{36}(n))| + \sqrt{30}|E_7(B_{36}(n))| + 6|E_8(B_{36}(n))|.$$

Then

$$R_{\frac{1}{2}}(G) = 8 + 8\sqrt{5}(-1 + n) + 46n + (6\sqrt{2} + 8\sqrt{3})(2 + n) + 16\sqrt{6}(1 + 2n) \\ + 6\sqrt{30}(-1 + 4n) + 18(-3 + 7n).$$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = -1$

$$R_{-1}(G) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} \frac{1}{\deg(u) \cdot \deg(v)}.$$

We have

$$R_{-1}(G) = \frac{1}{12}|E_1(B_{36}(n))| + \frac{1}{18}|E_2(B_{36}(n))| + \frac{1}{16}|E_3(B_{36}(n))| + \frac{1}{20}|E_4(B_{36}(n))| \\ + \frac{1}{24}|E_5(B_{36}(n))| + \frac{1}{25}|E_6(B_{36}(n))| + \frac{1}{30}|E_7(B_{36}(n))| + \frac{1}{36}|E_8(B_{36}(n))| \\ = \frac{1}{1800}(1255 + 5732n).$$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = -\frac{1}{2}$

$$R_{-\frac{1}{2}}(G) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} \frac{1}{\sqrt{\deg(u) \cdot \deg(v)}}.$$

Thus

$$R_{-\frac{1}{2}}(G) = \frac{\sqrt{3}}{6}|E_1(B_{36}(n))| + \frac{\sqrt{2}}{6}|E_2(B_{36}(n))| + \frac{1}{4}|E_3(B_{36}(n))| + \frac{\sqrt{5}}{10}|E_4(B_{36}(n))| \\ + \frac{\sqrt{6}}{12}|E_5(B_{36}(n))| + \frac{1}{5}|E_6(B_{36}(n))| + \frac{\sqrt{30}}{30}|E_7(B_{36}(n))| + \frac{1}{6}|E_8(B_{36}(n))| \\ = \frac{1}{30}(-30 + 20\sqrt{2} + 40\sqrt{3} - 12\sqrt{5} + 20\sqrt{6} - 6\sqrt{30} \\ + (171 + 10\sqrt{2} + 20\sqrt{3} + 12\sqrt{3} + 40\sqrt{6} + 24\sqrt{30})n).$$

□

In the following theorem, we compute first Zagreb index of borophene chain  $B_{36}(n)$ .

**Theorem 2.6.** For borophene chain  $G \cong B_{36}(n)$  for  $n \geq 2$ . Then

$$M_1(B_{36}(n)) = -22 + 850n.$$

*Proof.* Let  $G$  be the borophene chain  $B_{36}(n)$ . By using edge partition from Table 3, the result follows. From (4) we have

$$M_1(B_{36}(n)) = \sum_{uv \in E(G)} (deg(u) + deg(v)) = \sum_{j=1}^8 \sum_{uv \in E_j(G)} (deg(u) + deg(v)).$$

Then we have

$$M_1(B_{36}(n)) = 7|E_1(B_{36}(n))| + 9|E_2(B_{36}(n))| + 8|E_3(B_{36}(n))| + 9|E_4(B_{36}(n))| \\ + 10|E_5(B_{36}(n))| + 10|E_6(B_{36}(n))| + 11|E_7(B_{36}(n))| + 12|E_8(B_{36}(n))|.$$

By doing some calculation, we get  $M_1(B_{36}(n)) = -22 + 850n$ . □

Now, we compute  $ABC$  and  $GA$  indices of borophene chain  $B_{36}(n)$ .

**Theorem 2.7.** Let  $G \cong B_{36}(n)$  be the borophene chain, for  $n \geq 2$ , then

$$ABC(G) = \frac{1}{60}(24\sqrt{35}(-1+n) + 144\sqrt{2}n + (20\sqrt{14} + 40\sqrt{15})(2+n) \\ + (160\sqrt{3} + 30\sqrt{6})(1+2n) + 36\sqrt{30}(-1+4n) + 30\sqrt{10}(-3+7n)), \\ GA(G) = -7 + \frac{16}{9}\sqrt{5}(-1+n) + 31n + (\frac{48\sqrt{3} + 28\sqrt{2}}{21})(2+n) \\ + \frac{16}{5}\sqrt{6}(1+2n) + \frac{12}{11}\sqrt{30}(-1+4n).$$

*Proof.* By using edge partition given in Table 3, we get the result. From (5) it follows that

$$ABC(G) = \sum_{uv \in E(G)} \sqrt{\frac{deg(u) + deg(v) - 2}{deg(u) \cdot deg(v)}} = \sum_{j=1}^8 \sum_{uv \in E_j(G)} \sqrt{\frac{deg(u) + deg(v) - 2}{deg(u) \cdot deg(v)}}.$$

Then, we have

$$ABC(G) = \frac{\sqrt{15}}{6}|E_1(B_{36}(n))| + \frac{\sqrt{14}}{6}|E_2(B_{36}(n))| + \frac{\sqrt{6}}{4}|E_3(B_{36}(n))| \\ + \frac{\sqrt{35}}{10}|E_4(B_{36}(n))| + \frac{\sqrt{3}}{3}|E_5(B_{36}(n))| + 2\frac{\sqrt{2}}{5}|E_6(B_{36}(n))| \\ + \frac{\sqrt{30}}{10}|E_7(B_{36}(n))| + \frac{\sqrt{10}}{6}|E_8(B_{36}(n))|.$$

By doing some calculation, we get

$$ABC(G) = \frac{1}{60}(24\sqrt{35}(-1+n) + 144\sqrt{2}n + (20\sqrt{14} + 40\sqrt{15})(2+n) + (160\sqrt{3} + 30\sqrt{6})(1+2n) + 36\sqrt{30}(-1+4n) + 30\sqrt{10}(-3+7n)).$$

from (6) we get

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\deg(u)\deg(v)}}{(\deg(u) + \deg(v))} = \sum_{j=1}^8 \sum_{uv \in E_j(G)} \frac{2\sqrt{\deg(u)\deg(v)}}{(\deg(u) + \deg(v))}.$$

By doing some calculation, we get

$$GA(G) = 4\frac{\sqrt{3}}{7}|E_1(B_{36}(n))| + 2\frac{2}{3}|E_2(B_{36}(n))| + |E_3(B_{36}(n))| + 4\frac{\sqrt{5}}{9}|E_4(B_{36}(n))| + 2\frac{\sqrt{6}}{5}|E_5(B_{36}(n))| + |E_6(B_{36}(n))| + 2\frac{\sqrt{30}}{11}|E_7(B_{36}(n))| + |E_8(B_{36}(n))|.$$

We have

$$GA(G) = -7 + \frac{16}{9}\sqrt{5}(-1+n) + 31n + (\frac{48\sqrt{3} + 28\sqrt{2}}{21})(2+n) + \frac{16}{5}\sqrt{6}(1+2n) + \frac{12}{11}\sqrt{30}(-1+4n).$$

□

Now, we compute  $ABC_4$  and  $GA_5$  indices of borophene chain  $B_{36}(n)$ . Let us consider an edge partition based on degree sum of neighbors of end vertices. Then the edge set  $E(B_{36}(n))$  can be divided into twenty four edge partitions  $E_j(B_{36}(n)), 9 \leq j \leq 28$ , where the edge partition  $E_9(B_{36}(n))$  contains  $4n + 8$  edges  $uv$  with  $S_u = 14$  and  $S_v = 19$ , the edge partition  $E_{10}(B_{36}(n))$  contains  $2n + 4$  edges  $uv$  with  $S_u = 14$  and  $S_v = 28$ , the edge partition  $E_{11}(B_{36}(n))$  contains 6 edges  $uv$  with  $S_u = S_v = 19$ , the edge partition  $E_{12}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 19$  and  $S_v = 20$ , the edge partition  $E_{13}(B_{36}(n))$  contains  $4n + 8$  edges  $uv$  with  $S_u = 19$  and  $S_v = 28$ , the edge partition  $E_{14}(B_{36}(n))$  contains  $4n + 8$  edges  $uv$  with  $S_u = 19$  and  $S_v = 30$ , the edge partition  $E_{15}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 20$  and  $S_v = 26$ ,  $E_{16}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 20$  and  $S_v = 30$ ,  $E_{17}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 20$  and  $S_v = 31$ ,  $E_{18}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 26$  and  $S_v = 31$ ,  $E_{19}(B_{36}(n))$  contains  $2n - 2$  edges  $uv$  with  $S_u = 26$  and  $S_v = 35$ ,  $E_{20}(B_{36}(n))$  contains  $8n + 4$  edges  $uv$  with  $S_u = S_v = 28$ ,  $E_{21}(B_{36}(n))$  contains  $12n + 12$  edges  $uv$  with  $S_u = 28$  and  $S_v = 30$ ,  $E_{22}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 28$  and  $S_v = 31$ ,  $E_{23}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 28$  and  $S_v = 34$ ,  $E_{24}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 30$  and  $S_v = 31$ ,  $E_{25}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 31$  and  $S_v = 34$ ,  $E_{26}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 31$  and  $S_v = 35$ ,  $E_{27}(B_{36}(n))$  contains  $4n - 4$  edges  $uv$  with  $S_u = 34$  and  $S_v = 35$  and  $E_{28}(B_{36}(n))$  contains  $n - 1$  edges  $uv$  with  $S_u = S_v = 35$ .

**Theorem 2.8.** Let  $G \cong B_{36}(n)$  be the borophene chain, for  $n \geq 2$ , then

$$\begin{aligned}
 ABC_4(G) &= \frac{36}{19} + (6\sqrt{\frac{14}{527}} + 2\sqrt{\frac{134}{595}} + 2\sqrt{\frac{30}{119}} + 2\sqrt{\frac{118}{465}} + \sqrt{\frac{118}{455}} + 2\sqrt{\frac{57}{217}} \\
 &\quad + 2\sqrt{\frac{110}{403}} + 2\sqrt{\frac{22}{65}} + 2\sqrt{\frac{37}{95}} + \frac{4}{3}\sqrt{2} + \frac{2}{35}\sqrt{17} + \frac{14}{\sqrt{155}} + \frac{32}{\sqrt{1085}}) \\
 &\quad + 4\sqrt{\frac{3}{5}}(1+n) + (6\sqrt{\frac{5}{133}} + 2\sqrt{\frac{94}{285}} + 2\sqrt{\frac{62}{133}} + \frac{2}{7}\sqrt{5})(2+n) + \frac{3}{7}\sqrt{6}(1+2n), \\
 GA_5(G) &= 9 + (\frac{8}{5}\sqrt{6} + \frac{16}{39}\sqrt{95} + \frac{8}{23}\sqrt{130} + \frac{16}{51}\sqrt{155} + \frac{16}{59}\sqrt{217} + \frac{8}{31}\sqrt{238} + \frac{8}{57}\sqrt{806} \\
 &\quad + \frac{4}{61}\sqrt{910} + \frac{8}{61}\sqrt{930} + \frac{8}{65}\sqrt{1054} + \frac{4}{33}\sqrt{1085} + \frac{8}{69}\sqrt{1190})(-1+n) + 9n \\
 &\quad + \frac{24}{29}\sqrt{210}(1+n) + (\frac{4}{3}\sqrt{2} + \frac{16}{47}\sqrt{133} + \frac{8}{33}\sqrt{266} + \frac{16}{49}\sqrt{570})(2+n).
 \end{aligned}$$

*Proof.* By using edge partition given in Table 4, we get the result. From (7) it follows that

$$ABC_4(G) = \sum_{uv \in E(G)} \sqrt{\frac{S_u + S_v - 2}{S_u S_v}} = \sum_{j=9}^{28} \sum_{uv \in E_j(G)} \sqrt{\frac{S_u + S_v - 2}{S_u S_v}}.$$

Then, we have

$$\begin{aligned}
 ABC_4(G) &= \sqrt{\frac{31}{266}}|E_9(B_{36}(n))| + \frac{1}{7}\sqrt{5}|E_{10}(B_{36}(n))| + \frac{6}{19}|E_{11}(B_{36}(n))| \\
 &\quad + \frac{3}{2}\sqrt{\frac{5}{133}}|E_{13}(B_{36}(n))| + \sqrt{\frac{47}{570}}|E_{14}(B_{36}(n))| + \sqrt{\frac{15}{130}}|E_{14}(B_{36}(n))| \\
 &\quad + \frac{1}{3}\sqrt{2}|E_{16}(B_{36}(n))| + \frac{7}{2\sqrt{155}}|E_{17}(B_{36}(n))| + \sqrt{\frac{55}{806}}|E_{18}(B_{36}(n))| \\
 &\quad + \sqrt{\frac{59}{910}}|E_{19}(B_{36}(n))| + \frac{3}{28}\sqrt{6}|E_{20}(B_{36}(n))| + \frac{\sqrt{15}}{15}|E_{21}(B_{36}(n))| \\
 &\quad + \frac{1}{2}\sqrt{\frac{57}{217}}|E_{22}(B_{36}(n))| + \sqrt{\frac{15}{238}}|E_{23}(B_{36}(n))| + \sqrt{\frac{59}{930}}|E_{24}(B_{36}(n))| \\
 &\quad + 3\sqrt{\frac{7}{1054}}|E_{25}(B_{36}(n))| + \frac{8}{\sqrt{1085}}|E_{26}(B_{36}(n))| + \sqrt{\frac{67}{1190}}|E_{27}(B_{36}(n))| \\
 &\quad + \frac{2}{35}\sqrt{17}|E_{28}(B_{36}(n))| + \frac{1}{2}\sqrt{\frac{37}{95}}|E_{12}(B_{36}(n))|.
 \end{aligned}$$

Thus, we have

$$\begin{aligned}
 ABC_4(G) &= \frac{36}{19} + (6\sqrt{\frac{14}{527}} + 2\sqrt{\frac{134}{595}} + 2\sqrt{\frac{30}{119}} + 2\sqrt{\frac{118}{465}} + \sqrt{\frac{118}{455}} + 2\sqrt{\frac{57}{217}} + 2\sqrt{\frac{110}{403}} \\
 &\quad + 2\sqrt{\frac{22}{65}} + 2\sqrt{\frac{37}{95}} + \frac{4}{3}\sqrt{2} + \frac{2}{35}\sqrt{17} + \frac{14}{\sqrt{155}} + \frac{32}{\sqrt{1085}}) + 4\sqrt{\frac{3}{5}}(1+n) \\
 &\quad + (6\sqrt{\frac{5}{133}} + 2\sqrt{\frac{94}{285}} + 2\sqrt{\frac{62}{133}} + \frac{2}{7}\sqrt{5})(2+n) + \frac{3}{7}\sqrt{6}(1+2n),
 \end{aligned}$$



Table 4. Edge partition of borophene chain  $B_{36}(n)$  based on degrees sum of end vertices of each edge.

$(S_u, S_v), uv \in E(G)$	Number of edges	$(S_u, S_v), uv \in E(G)$	Number of edges
(14, 19)	$4n + 8$	(26, 35)	$2n - 2$
(14, 28)	$2n + 4$	(28, 28)	$8n + 4$
(19, 19)	6	(28, 30)	$12n + 12$
(19, 20)	$4n - 4$	(28, 31)	$4n - 4$
(19, 28)	$4n + 8$	(28, 34)	$4n - 4$
(19, 30)	$4n + 8$	(30, 31)	$4n - 4$
(20, 26)	$4n - 4$	(31, 34)	$4n - 4$
(20, 30)	$4n - 4$	(31, 35)	$4n - 4$
(20, 31)	$4n - 4$	(34, 35)	$4n - 4$
(26, 31)	$4n - 4$	(35, 35)	$n - 1$

and from (8) we get

$$GA_5(G) = \sum_{uv \in E(G)} \frac{2\sqrt{S_u S_v}}{(S_u + S_v)} = \sum_{j=9}^{28} \sum_{uv \in E_j(G)} \frac{2\sqrt{S_u S_v}}{(S_u + S_v)}.$$

Then, we have

$$\begin{aligned} GA_5(G) &= \frac{2}{33} \sqrt{266} |E_9(B_{36}(n))| + \frac{2}{3} \sqrt{2} |E_{10}(B_{36}(n))| + |E_{11}(B_{36}(n))| \\ &+ \frac{4}{39} \sqrt{95} |E_{12}(B_{36}(n))| + \frac{4}{47} \sqrt{133} |E_{13}(B_{36}(n))| + \frac{4}{49} \sqrt{570} |E_{14}(B_{36}(n))| \\ &+ \frac{2}{23} \sqrt{130} |E_{15}(B_{36}(n))| + \frac{2}{5} \sqrt{6} |E_{16}(B_{36}(n))| + \frac{4}{51} \sqrt{155} |E_{17}(B_{36}(n))| \\ &+ \frac{2}{57} \sqrt{806} |E_{18}(B_{36}(n))| + \frac{2}{61} \sqrt{910} |E_{19}(B_{36}(n))| + |E_{20}(B_{36}(n))| \\ &+ \frac{2}{29} \sqrt{210} |E_{21}(B_{36}(n))| + \frac{4}{59} \sqrt{214} |E_{22}(B_{36}(n))| + \frac{2}{31} \sqrt{238} |E_{23}(B_{36}(n))| \\ &+ \frac{2}{61} \sqrt{930} |E_{24}(B_{36}(n))| + \frac{2}{65} \sqrt{1054} |E_{25}(B_{36}(n))| + \frac{1}{33} \sqrt{1085} |E_{26}(B_{36}(n))| \\ &+ \frac{2}{69} \sqrt{1190} |E_{27}(B_{36}(n))| + |E_{28}(B_{36}(n))| \\ &= 9 + \left(\frac{8}{5} \sqrt{6} + \frac{16}{39} \sqrt{95} + \frac{8}{23} \sqrt{130} + \frac{16}{51} \sqrt{155} + \frac{16}{59} \sqrt{217} + \frac{8}{31} \sqrt{238} + \frac{8}{57} \sqrt{806}\right. \\ &+ \frac{4}{61} \sqrt{910} + \frac{8}{61} \sqrt{930} + \frac{8}{65} \sqrt{1054} + \frac{4}{33} \sqrt{1085} + \frac{8}{69} \sqrt{1190} \Big) (-1 + n) + 9n \\ &+ \frac{24}{29} \sqrt{210} (1 + n) + \left(\frac{4}{3} \sqrt{2} + \frac{16}{47} \sqrt{133} + \frac{8}{33} \sqrt{266} + \frac{16}{49} \sqrt{570}\right) (2 + n). \end{aligned}$$

□

The melem (2, 5, 8-triamino-tri-s-triazine)  $C_6N_7(NH_2)_3$  chain nanotube. The melem was obtained as a crystalline powder by thermal treatment of different less condensed C – N – H

compounds (e.g., melamine  $C_3N_3(NH_2)_3$ , dicyandiamide  $H_4C_2N_4$ , ammonium dicyanamide  $NH_4[N(CN)_2]$ , or cyanamide  $H_2CN_2$ , respectively) at temperatures up to  $450^\circ C$  in sealed glass ampules. The vertices and edges in melem chain are  $18n + 4$  and  $21n + 3$  respectively.

Now we compute Randić  $R_\alpha(G)$  with  $\alpha = \{1, -1, \frac{1}{2}, -\frac{1}{2}\}$ ,  $ABC$ ,  $GA$ ,  $ABC_4$  and  $GA_5$  indices for melem chain  $MC(n)$  nanotube.

**Theorem 2.9.** Consider the melem chain  $MC(n)$  for  $n \in \mathbb{N}$ . Then

$$R_\alpha(MC(n)) = \begin{cases} 135n + 9, & \alpha = 1; \\ 3(6n + 4\sqrt{6}n + \sqrt{3}(1 + n)), & \alpha = \frac{1}{2}; \\ 1 + \frac{11n}{3}, & \alpha = -1; \\ \sqrt{3} + (2 + \sqrt{3} + 2\sqrt{6})n, & \alpha = -\frac{1}{2}. \end{cases}$$

*Proof.* Let  $G$  be the melem chain. The melem chain  $MC(n)$  has  $3n + 3$  vertices of degree 1,  $6n$  vertices of degree 2, and  $9n + 1$  vertices of degree 3. The edge set of  $MC(n)$  is divided into three partitions based on the degree of end vertices. The first edge partition  $E_1(MC(n))$  contains  $3n + 3$  edges  $uv$ , where  $deg(u) = 1$  and  $deg(v) = 3$ . The second edge partition  $E_2(MC(n))$  contains  $12n$  edges  $uv$ , where  $deg(u) = 2$  and  $deg(v) = 3$ . The third edge partition  $E_3(MC(n))$  contains  $6n$  edges  $uv$ , where  $deg(u) = deg(v) = 3$ . Table 5 shows such an edge partition of  $MC(n)$ . Thus from (3) it follows that

$$R_\alpha(G) = \sum_{uv \in E(G)} (deg(u)deg(v))^\alpha.$$

Now we apply the formula of  $R_\alpha(G)$  for  $\alpha = 1$

$$R_1(G) = \sum_{j=1}^3 \sum_{uv \in E_j(G)} deg(u)deg(v).$$

By using edge partition given in Table 5, we get

$$R_1(G) = 3|E_1(MC(n))| + 6|E_2(MC(n))| + 9|E_3(MC(n))| = 135n + 9.$$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = \frac{1}{2}$

$$R_{\frac{1}{2}}(G) = \sum_{j=1}^3 \sum_{uv \in E_j(G)} \sqrt{deg(u) \cdot deg(v)}.$$

By using edge partition given in Table 5, we get

$$\begin{aligned} R_{\frac{1}{2}}(G) &= \sqrt{3}|E_1(MC(n))| + \sqrt{6}|E_2(MC(n))| + 3|E_3(MC(n))| \\ &= 3(6n + 4\sqrt{6}n + \sqrt{3}(1 + n)). \end{aligned}$$

Table 5. Edge partition of melem chain  $MC(n)$  based on degrees of end vertices of each edge.

$(d_u, d_v), uv \in E(G)$	Number of edges
(1,3)	$3n + 3$
(2,3)	$12n$
(3,3)	$6n$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = -1$

$$\begin{aligned} R_{-1}(G) &= \sum_{j=1}^3 \sum_{uv \in E_j(G)} \frac{1}{deg(u) \cdot deg(v)} \\ &= \frac{1}{3}|E_1(MC(n))| + \frac{1}{6}|E_2(MC(n))| + \frac{1}{9}|E_3(MC(n))| \\ &= 1 + \frac{11n}{3}. \end{aligned}$$

We apply the formula of  $R_\alpha(G)$  for  $\alpha = -\frac{1}{2}$

$$\begin{aligned} R_{-\frac{1}{2}}(G) &= \sum_{j=1}^3 \sum_{uv \in E_j(G)} \frac{1}{\sqrt{deg(u) \cdot deg(v)}} \\ &= \frac{1}{\sqrt{3}}|E_1(MC(n))| + \frac{1}{\sqrt{6}}|E_2(MC(n))| + \frac{1}{3}|E_3(MC(n))| \\ &= \sqrt{3} + (2 + \sqrt{3} + 2\sqrt{6})n. \end{aligned}$$

□

In the following theorem, we compute first Zagreb index of melem chain  $MC(n)$ .

**Theorem 2.10.** For melem chain  $G \cong MC(n)$  for  $n \in \mathbb{N}$ . Then

$$M_1(MC(n)) = 12(1 + 9n).$$

*Proof.* Let  $G$  be the borophene chain  $B_{36}(n)$ . By using edge partition from Table 5, the result follows. From (4) we have

$$\begin{aligned} M_1(MC(n)) &= \sum_{uv \in E(G)} (deg(u) + deg(v)) \\ &= \sum_{j=1}^3 \sum_{uv \in E_j(G)} (deg(u) + deg(v)) \\ &= 4|E_1(MC(n))| + 5|E_2(MC(n))| + 6|E_3(MC(n))|. \end{aligned}$$

By doing some calculation, we get  $M_1(MC(n)) = 12(1 + 9n)$ .

□

Now, we compute  $ABC$  and  $GA$  indices of melem chain  $MC(n)$ .

**Theorem 2.11.** Let  $G \cong MC(n)$  be the melem chain, for  $n \in \mathbf{N}$ , then

$$ABC(G) = \sqrt{6} + (4 + 6\sqrt{2} + \sqrt{6})n,$$

$$GA(G) = 6n + \frac{24\sqrt{6}}{5}n + \frac{3\sqrt{3}}{2}(1 + n).$$

*Proof.* By using edge partition given in Table 5, we get the result. From (5) it follows that

$$\begin{aligned} ABC(G) &= \sum_{uv \in E(G)} \sqrt{\frac{\deg(u) + \deg(v) - 2}{\deg(u) \cdot \deg(v)}} \\ &= \sum_{j=1}^3 \sum_{uv \in E_j(G)} \sqrt{\frac{\deg(u) + \deg(v) - 2}{\deg(u) \cdot \deg(v)}} \\ &= \sqrt{\frac{2}{3}}|E_1(MC(n))| + \frac{1}{\sqrt{2}}|E_2(MC(n))| + \frac{2}{3}|E_3(MC(n))|. \end{aligned}$$

By doing some calculation, we get  $ABC(G) = \sqrt{6} + (4 + 6\sqrt{2} + \sqrt{6})n$ , from (6) we get

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\deg(u)\deg(v)}}{(\deg(u) + \deg(v))} = \sum_{j=1}^3 \sum_{uv \in E_j(G)} \frac{2\sqrt{\deg(u)\deg(v)}}{(\deg(u) + \deg(v))}.$$

By doing some calculation, we get

$$\begin{aligned} GA(G) &= \frac{\sqrt{3}}{2}|E_1(MC(n))| + \frac{2\sqrt{6}}{5}|E_2(MC(n))| + |E_3(MC(n))|, \\ &= 6n + \frac{24\sqrt{6}}{5}n + \frac{3\sqrt{3}}{2}(1 + n). \end{aligned}$$

□

Now, we compute  $ABC_4$  and  $GA_5$  indices of melem chain  $MC(n)$ . Let us consider an edge partition based on degree sum of neighbors of end vertices. Then the edge set  $E(MC(n))$  can be divided into six edge partitions  $E_j(MC(n)), 4 \leq j \leq 9$ , where the edge partition  $E_4(MC(n))$  contains  $2n + 4$  edges  $uv$  with  $S_u = 3$  and  $S_v = 5$ , the edge partition  $E_5(MC(n))$  contains  $n - 1$  edges  $uv$  with  $S_u = 3$  and  $S_v = 7$ , the edge partition  $E_6(MC(n))$  contains  $n + 2$  edges  $uv$  with  $S_u = 5$  and  $S_v = 7$ , the edge partition  $E_7(MC(n))$  contains  $12n$  edges  $uv$  with  $S_u = 6$  and  $S_v = 7$ , the edge partition  $E_8(MC(n))$  contains  $2n - 2$  edges  $uv$  with  $S_u = S_v = 7$  and the edge partition  $E_9(MC(n))$  contains  $3n$  edges  $uv$  with  $S_u = 7$  and  $S_v = 9$ .

**Theorem 2.12.** Let  $G \cong MC(n)$  be the melem chain, for  $n \geq 2$ , then

$$\begin{aligned} ABC_4(G) &= (2\sqrt{\frac{2}{21}} + \frac{4\sqrt{3}}{7})(-1 + n) + (\sqrt{2} + 2\sqrt{\frac{66}{7}})n + (\sqrt{\frac{2}{7}} + 2\sqrt{\frac{2}{5}})(2 + n), \\ GA_5(G) &= -2 + \frac{1}{5}\sqrt{21}(-1 + n) + (2 + \frac{9\sqrt{7}}{8} + \frac{24\sqrt{42}}{13})n \\ &\quad + (\frac{1}{2}\sqrt{15} + \frac{1}{6}\sqrt{35})(2 + n). \end{aligned}$$

Table 6. Edge partition of Melem chain  $MC(n)$  based on degrees sum of end vertices of each edge.

$(S_u, S_v), uv \in E(G)$	Number of edges
(3,5)	$2n + 4$
(3,7)	$n - 1$
(5,7)	$n + 2$
(6,7)	$12n$
(7,7)	$2n - 2$
(7,9)	$3n$

*Proof.* By using edge partition given in Table 6, we get the result. From (7) it follows that

$$\begin{aligned}
 ABC_4(G) &= \sum_{uv \in E(G)} \sqrt{\frac{S_u + S_v - 2}{S_u S_v}} = \sum_{j=4}^9 \sum_{uv \in E_j(G)} \sqrt{\frac{S_u + S_v - 2}{S_u S_v}} \\
 &= \sqrt{\frac{2}{5}} |E_4(MC(n))| + \frac{2\sqrt{2}}{\sqrt{21}} |E_5(MC(n))| + \sqrt{\frac{2}{7}} |E_6(MC(n))| \\
 &+ \sqrt{\frac{11}{42}} |E_7(MC(n))| + \frac{2\sqrt{3}}{7} |E_8(MC(n))| + \frac{\sqrt{2}}{3} |E_9(MC(n))| \\
 &= (2\sqrt{\frac{2}{21}} + \frac{4\sqrt{3}}{7})(-1 + n) + (\sqrt{2} + 2\sqrt{\frac{66}{7}})n + (\sqrt{\frac{2}{7}} + 2\sqrt{\frac{2}{5}})(2 + n),
 \end{aligned}$$

and from (8) we get

$$\begin{aligned}
 GA_5(G) &= \sum_{uv \in E(G)} \frac{2\sqrt{S_u S_v}}{(S_u + S_v)} = \sum_{j=4}^9 \sum_{uv \in E_j(G)} \frac{2\sqrt{S_u S_v}}{(S_u + S_v)} \\
 &= \frac{\sqrt{15}}{4} |E_4(MC(n))| + \frac{\sqrt{21}}{5} |E_5(MC(n))| + \frac{\sqrt{35}}{6} |E_6(MC(n))| \\
 &+ \frac{2\sqrt{42}}{13} |E_7(MC(n))| + |E_8(MC(n))| + \frac{\sqrt{63}}{8} |E_9(MC(n))| \\
 &= -2 + \frac{1}{5}\sqrt{21}(-1 + n) + (2 + \frac{9\sqrt{7}}{8} + \frac{24\sqrt{42}}{13})n + (\frac{1}{2}\sqrt{15} + \frac{1}{6}\sqrt{35})(2 + n).
 \end{aligned}$$

□

### 3 Conclusion

In this paper, certain degree based topological indices, namely general Randić index, atomic-bond connectivity index ( $ABC$ ), geometric-arithmetic index ( $GA$ ) and first Zagreb index for boron triangular sheet  $BTS(m, n)$ , borophene chain of  $B_{36}(n)$  and melem chain  $MC(n)$  were studied for the first time and analytical closed formulas for these nanostructure were determined which will help the people working in chemical science to understand and explore the underlying topologies of these nanostructures.

## 4 Acknowledgment

The authors are very grateful to the referees for their careful reading with corrections and useful comments, which improved this work very much.

## References

- [1] H. Ali, A. Q. Baig, M. K. Shafiq, On topological properties of hierarchical interconnection networks, *J. Appl. Math. Comput.* 55 (1-2) (2017) 313–334.
- [2] M. Bača, J. Horváthová, M. Mokrišová, A. Suhányiová, On topological indices of fullerenes, *Appl. Math. Comput.* 251 (2015) 154–161.
- [3] A. Q. Baig, M. Imran, H. Ali, Computing Omega, Sadhana and PI polynomials of benzoid carbon nanotubes, *OAM-RC.* 9 (2015) 248–255.
- [4] A. Q. Baig, M. Imran, H. Ali, On Topological Indices of Poly Oxide, Poly Silicate, DOX and DSL Networks, *Can. J. Chem.* 93 (2015) 1–10.
- [5] M. Deza, P. W. Fowler, A. Rassat, K. M. Rogers, Fullerenes as tiling of surfaces, *J. Chem. Inf. Comput. Sci.* 40 (2000) 550–558.
- [6] M. V. Diudea, I. Gutman, J. Lorentz, *Molecular Topology*, Nova, Huntington, 2001.
- [7] E. Estrada, L. Torres, L. Rodríguez, I. Gutman, An atom-bond connectivity index: Modelling the enthalpy of formation of alkanes, *Indian J. Chem.* 37A (1998) 849–855.
- [8] M. Ghorbani, M. A. Hosseinzadeh, Computing  $ABC_4$  index of nanostar dendrimers, *Optoelectron. Adv. Mater. Rapid Commun.* 4 (2010) 1419–1422.
- [9] A. Graovac, M. Ghorbani, M. A. Hosseinzadeh, Computing fifth geometric-arithmetic index for nanostar dendrimers, *J. Math. Nanosci.* 1 (2011) 33–42.
- [10] I. Gutman, O. E. Polansky, *Mathematical concepts in organic chemistry*, Springer-Verlag, New York, 1986.
- [11] S. Hayat, M. Imran, Computation of certain topological indices of nanotubes, *J. Comput. Theor. Nanosci.* 12 (2015) 70–76.
- [12] S. Hayat, M. Imran, Computation of topological indices of certain networks, *Appl. Math. Comput.* 240 (2014) 213–228.
- [13] M. Imran, A. Q. Baig, H. Ali, On topological properties of dominating David derived graphs, *Can. J. Chem.* 94 (2016) 137–148.
- [14] M. Imran, A. Q. Baig, H. Ali, On molecular topological properties of hex-derived graphs, *J. Chemometrics*, 30 (2016) 121–129.
- [15] M. Imran, A. Q. Baig, H. Ali, S. U. Rehman, On topological properties of poly honeycomb graphs, *Period Math Hung.* 73 (2016) 100–119.
- [16] A. Iranmanesh, M. Zeraatkar, Computing GA index for some nanotubes, *Optoelectron. Adv. Mater. Rapid Commun.* 4 (2010) 1852–1855.
- [17] W. Lin, J. Chen, Q. Chen, T. Gao, X. Lin, B. Cai, Fast computer search for trees with minimal ABC index based on tree degree sequences, *MATCH Commun. Math. Comput. Chem.* 72 (2014) 699–708.
- [18] P. D. Manuel, M. I. Abd-El-Barr, I. Rajasingh, B. Rajan, An efficient representation of Benes networks and its applications, *J. Discrete Algorithms*, 6 (2008) 11–19.
- [19] J. L. Palacios, A resistive upper bound for the ABC index, *MATCH Commun. Math. Comput. Chem.* 72 (2014) 709–713.
- [20] M. Randić, On Characterization of molecular branching, *J. Amer. Chem. Soc.* 97 (1975) 6609–6615.
- [21] F. Simonraj, A. George, Embedding of poly honeycomb networks and the metric dimension of star of david network, *GRAPH-HOC*, 4 (2012) 11–28.
- [22] F. Simonraj, A. George, Topological Properties of few Poly Oxide, Poly Silicate, DOX and DSL Networks, *International Journal of Future Computer and Communication*, 2 (2013) 90–95.
- [23] D. Vukičević B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges, *J. Math. Chem.* 46 (2009) 1369–1376.
- [24] H. Wiener, Structural determination of paraffin boiling points, *J. Amer. Chem. Soc.* 69 (1947) 17–20.